# **PSRI INSTITUTE OF ORTHOPAEDICS & JOINT REPLACEMENT**



DR. PRAKASH P. KOTWAL

MBBS, MS, (Ortho), FAMS, FIMSA CHAIRMAN Institute of Orthopaedics & Joint Replacement Formerly Professor & Head, Dept. Of Orthopaedics, AIIMS, New Delhi

Reg No: 12112 (DMC)

**PUSHPAWATI SINGHANIA** HOSPITAL & RESEARCH INSTITUTE

Mobile No: 9868129599

E-mail id : prakash.kotwal@gmail.com

Date: 17 08 202

ARVIND MISRA 454R/ more

FUC 9 TOTO WISH FUSION (E) FIX SNAW WISH (E) Dosy - 31-7-2021

Ward a Survey lin

Ward a Survey lin

John - BWRIST SPENT

GHAR - CONTINUE CARUUM 4 VIT D3

- CONTINUE B-COMPLEX.

- RIV AFTER 6 WEEKS X-RAY (2) WRIST ( CONEM) BY 17.67.204



15/9/2021

The Pain (toward Ulrar Side) (B) Mand.

Today +x over Lithe first

- BONE SLAN (to we port) - S. PTH

TAB TRO CONTINE 100 my X10 DAYS TAB SOMPRAZ D YON 1. 00 × 10 mg

( Sepre brandost)

TAB ME-12 00 1.00 × I MENTH. But

05/10/24 S. P14-36.2 (N) BINE SLAW 4) S/O RSD @ NAST

Av - TWO BEN157 4 20 mg S. C OD X 3 mw/m - tas cen 1. 00 × 3 months THE UPRISE D3 ONCE MINTHLY

TAB ME-12 00 1. 00 x 2 mws4

- allescateat

By 3/10/2021





#### **DEPARTMENT OF NUCLEAR MEDICINE AND PET-CT**

NAME	MR. ARVIND MISHRA	AGE/SEX	45 Y/M	DATE:	18.09.2021
REG NO	60057336	REF BY	DR. PRAKASH P. KOTWAL		

Clinical History:- C/o right wrist pain post total wrist fusion (31.07.2021).

### Procedure:-

### 99mTc MDP Three Phase Skeletal Scintigraphy with SPECT/CT

Sequential dynamic images of pelvic region were acquired in anterior and posterior views immediately after intravenous administration of Tc-99m MDP. Regional flow and blood pool images were acquired after 1 min followed by whole body blood pool images. Regional and whole body delayed images were acquired 3 hrs after tracer administration.

#### FINDINGS:

- Regional perfusion and blood pool images: Increased flow and blood pool in right wrist.
- Delayed regional and whole body images: Increased tracer uptake noted in right wrist.
- Physiological tracer uptake is noted in rest of the skeleton.
- Both the kidneys are well visualized and normal in position.

### **IMPRESSION:**

> Increased tracer activity in all phases of bone scan suggestive of RSD.

Dr. ASHUTOSH PARASHAR

MBBS, DNB (Nuclear Medicine)
PDCC (Co-relative Imaging)

Ex. Senior Resident (SGPGI)

Consultant-Nuclear Medicine & PET-CT

Disclaimer:

This is a professional opinion and not a definite diagnosis. Further clinico-pathological correlation is necessary. Please interpret accordingly. This report is not valid for medico legal purpose. For any typing related errors kindly report us within 15 days with original report of examination and supporting documents as proof.

## Shri Ram Murti Smarak Functional Imaging & Medical Centres e 1 12 10000

CP 2/3, Vishwas Khand -2, Near Flyover, Gomti Nagar, Lucknow-226010 Ph.: 0522-2308987-88 Mob.: 9458704154 Fax: 0522-2308986 E-mail: srmsfimc@srms.ac.in Website: srms.ac.in/fimc Working Hours: 8 a.m. to 9 p.m. (Mon-Sat), 9 am to 6 p.m. (Sunday/ Holidays) Diagnostic Services: PET-CT Scan, SPECT Gamma Dual Camera, 3T MRI 48 Channels, CT Scan 128 Slice, Digital Mammography Digital X-Ray, 4D USG Color Doppler, Pathology, Microbiology, Biochemistry, ECG, TMT, PFT